

Macroscopic Thermodynamical Witnesses of Quantum Entanglement

Časlav Brukner^{1,2,*} and Vlatko Vedral^{1,†}

¹*Optics Section, The Blackett Laboratory, Imperial College,
Prince Consort Road, London, SW7 2BW, United Kingdom*

²*Institut für Experimentalphysik, Universität Wien, Boltzmannngasse 5, A-1090 Wien, Austria*
(Dated: February 1, 2008)

We show that macroscopic thermodynamical properties - such as functions of internal energy and magnetization - can detect quantum entanglement in solids at nonzero temperatures in the thermodynamical limit. We identify the parameter regions (critical values of magnetic field and temperature) within which entanglement is witnessed by these thermodynamical quantities.

PACS numbers: 03.67.Hk, 03.65.Ta, 03.65.Ud

An entangled quantum system is impossible to describe by the states of its (local) constituents alone [1]. This fundamental feature of quantum mechanics can be manifested in a phenomenon known as quantum non-locality [2]. Besides being of fundamental interest, entanglement is considered as the crucial resource for quantum information processing [3], but its effects are not generally seen beyond the atomic scale and only in well controlled laboratory conditions. It is an intriguing and fascinating question to what extent entanglement may develop naturally in realistic complex systems and can affect their macroscopic properties.

Recently, extensive efforts have been made to understand theoretically and to quantify entanglement in solid state systems [4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19]. Within various models of interacting spins in arrays entanglement was found to be present at moderate nonzero temperatures – phenomenon known as *thermal entanglement* [4, 5] – and has been linked to the existence of critical phenomena [10, 11, 12, 13, 20]. Most of the studies, however, were limited to only small number of spin sites. On the other hand, in the thermodynamical limit (where the number N of spins tends to infinity) entanglement was predominately investigated for the systems in their ground states [9, 10, 11, 12, 13, 14].

Only very few results are known about a possibility of existence of entanglement and its properties in the thermal states in the thermodynamical limit. Entanglement as measured by concurrence [21] was shown to exist in this limit at nonzero temperatures in the transverse Ising model [10] and the antiferromagnetic Heisenberg model [7, 8] and an upper bound for the multipartite entanglement in the transverse Ising model was given [16]. Finally, it was suggested that macroscopic entanglement is possible even at high temperatures (as high as 160 Kelvin) in high-temperature superconductors [17] (see also Ref. [18]).

Thermodynamic laws are of very general validity, and they do not depend on the details of the interactions or type of the (microscopic) constituents of the system being studied. When the system is at thermal equilibrium under a certain temperature T , it is in a

thermal state $\rho = e^{-H/kT}/Z$, where $Z = \text{Tr}(e^{-H/kT})$ is the partition function, H is the Hamiltonian and k is the Boltzmann constant. The partition function is the central object of statistical physics from which all other thermodynamical quantities can be derived: e.g. internal energy $U = -(1/Z)(\partial Z/\partial\beta)$ or magnetization $M = -(1/Z\beta)(\partial Z/\partial B)$, where $\beta = 1/kT$. It is generally believed that although the partition function is determined by the eigenvalues of Hamiltonian only, detecting entanglement in solids requires in addition the knowledge of the energy eigenstates. This is in general a hard problem and origin of the main difficulties in the research on entanglement in solid state systems.

Recently, however, it was demonstrated experimentally that entanglement can affect macroscopic properties of solids, albeit at very low (critical) temperature (below 1 Kelvin) [19]. This extraordinary result opens up a possibility that purely quantum correlations between microscopic constituents of the solid may be detected by only a small number of macroscopic thermodynamical properties. In the similar spirit concurrence in the isotropic XXX Heisenberg model [7, 8] and an upper bound on entanglement in the transverse Ising model [16] was given in terms of internal energy.

In this Letter we show that thermodynamical macroscopic properties can serve as (multipartite) entanglement witnesses for the thermal states in the thermodynamical limit. Entanglement witnesses are observables which have positive expectation values for separable states and negative one for some, specific, entangled states [22]. We show that a function of internal energy and magnetization can detect entanglement in the thermal states in the (XX and XXX) Heisenberg models for a wide range of values of the external magnetic field and temperature. The critical temperature below which entanglement is present in the system is of the order of the coupling constant J (Experimental estimations of this constant gives the values as high as about 10 Kelvin [23], where J is measured in kT units. Compare with Ref. [17]).

We consider the linear (1D) spin chain in the Heisenberg model. The interaction is between nearest-neighbor

spins and the external field B is along z direction. The Hamiltonian is given by

$$H = - \sum_{i=1}^N (J_x \sigma_i^x \sigma_{i+1}^x + J_y \sigma_i^y \sigma_{i+1}^y + J_z \sigma_i^z \sigma_{i+1}^z) - B \sum_{i=1}^N \sigma_i^z, \quad (1)$$

where σ_i^x , σ_i^y , and σ_i^z are the Pauli spin operators for the i th spin. Throughout the paper we will consider special cases of this Hamiltonian – the isotropic XXX Heisenberg model with $J_x = J_y = J_z = J$ and isotropic XX Heisenberg model with $J_x = J_y = J$ and $J_z = 0$. The regimes $J > 0$ and $J < 0$ correspond to the antiferromagnetic and the ferromagnetic cases, respectively.

We will now derive a thermodynamical entanglement witness for a solid state system in thermal equilibrium. Using $U = \langle H \rangle$ and $M = \sum_{j=1}^N \langle \sigma_j^z \rangle$ we obtain

$$\frac{U + BM}{NJ} = -\frac{1}{N} \sum_{i=1}^N (\langle \sigma_i^x \sigma_{i+1}^x \rangle + \langle \sigma_i^y \sigma_{i+1}^y \rangle + \langle \sigma_i^z \sigma_{i+1}^z \rangle) \quad (2)$$

from Eq. (1). The right-hand of Eq. (2) is an entanglement witness as shown in Ref. [24]: for any separable state, that is, for any classical mixture of the products states: $\rho = \sum_k w_k \rho_k^1 \otimes \rho_k^2 \otimes \dots \otimes \rho_k^N$, one has

$$\frac{1}{N} \left| \sum_{i=1}^N (\langle \sigma_i^x \sigma_{i+1}^x \rangle + \langle \sigma_i^y \sigma_{i+1}^y \rangle + \langle \sigma_i^z \sigma_{i+1}^z \rangle) \right| \leq 1. \quad (3)$$

The proof is based on the fact that for any product state $\rho_k^1 \otimes \dots \otimes \rho_k^N$ and for every i one has $|\langle \sigma_i^x \sigma_{i+1}^x \rangle + \langle \sigma_i^y \sigma_{i+1}^y \rangle + \langle \sigma_i^z \sigma_{i+1}^z \rangle| = |\langle \sigma_i^x \rangle \langle \sigma_{i+1}^x \rangle + \langle \sigma_i^y \rangle \langle \sigma_{i+1}^y \rangle + \langle \sigma_i^z \rangle \langle \sigma_{i+1}^z \rangle| \leq \sqrt{\langle \sigma_i^x \rangle^2 + \langle \sigma_i^y \rangle^2 + \langle \sigma_i^z \rangle^2} \sqrt{\langle \sigma_{i+1}^x \rangle^2 + \langle \sigma_{i+1}^y \rangle^2 + \langle \sigma_{i+1}^z \rangle^2} \leq 1$. This is also valid for any convex sum of product states (separable states). The upper bound was found by using the Cauchy-Schwarz inequality and knowing that for any state $\langle \sigma^x \rangle^2 + \langle \sigma^y \rangle^2 + \langle \sigma^z \rangle^2 \leq 1$. It is important to note that the same proof can also be applied if one considers XX Heisenberg model. In this case one has

$$\frac{1}{N} \left| \sum_{i=1}^N (\langle \sigma_i^x \sigma_{i+1}^x \rangle + \langle \sigma_i^y \sigma_{i+1}^y \rangle) \right| \leq 1 \quad (4)$$

for any separable state.

We now give our thermodynamical entanglement witness: if, in the isotropic XXX or XX Heisenberg model, one has

$$\frac{|U + BM|}{N|J|} > 1, \quad (5)$$

then the solid state system is in an entangled state. The entanglement witness is physically equivalent to the exchange interaction energy or, equivalently, to the difference between the total (internal) energy U and the magnetic energy $-BM$. From the tracelessness of the Pauli

operators one can easily see that $\lim_{T \rightarrow \infty} U \rightarrow 0$. This means that the value of the internal energy as given by (5) should be defined relatively to the referent value of zero energy in the limit of high temperatures.

We now give an explicit example of a state that violates Ineqs. (3) and (4). This then completes the proof that expression (5) is indeed an entanglement witness and not just a bound that is trivially satisfied by any quantum state. As an example of such a state we take the ground state of the antiferromagnetic isotropic XXX Heisenberg model with zero magnetic field. The energy of this state was found to be [9, 25]: $|E_0/JN| = 1/N |\sum_{i=1}^N (\langle \sigma_i^x \sigma_{i+1}^x \rangle_0 + \langle \sigma_i^y \sigma_{i+1}^y \rangle_0 + \langle \sigma_i^z \sigma_{i+1}^z \rangle_0)| = 1.773 > 1$, where the index “0” denotes that the mean values are taken for the ground state. Furthermore, due to the symmetry of the XXX Heisenberg Hamiltonian one has $E_0/(3NJ) = \langle \sigma_i^x \sigma_{i+1}^x \rangle_0 = \langle \sigma_i^y \sigma_{i+1}^y \rangle_0 = \langle \sigma_i^z \sigma_{i+1}^z \rangle_0 = -1.773/3$ for every i . This implies that $1/N |\sum_{i=1}^N (\langle \sigma_i^x \sigma_{i+1}^x \rangle_0 + \langle \sigma_i^y \sigma_{i+1}^y \rangle_0)| = 1.182 > 1$. Therefore, Ineq. (5) is an entanglement witness for the solid state systems described by XXX or XX Heisenberg interaction.

We will now discuss various concrete models of spin interaction of which some are exactly solvable and for which dependence of internal energy U and magnetization M on temperature T and magnetic field B are known. This will help us to determine the parameter regions of T and B within which one has entanglement in the solids.

We first consider *XXX Heisenberg model with no magnetic field* ($J_x = J_y = J_z = J$ and $B = 0$). Since there is no magnetic field, symmetry requires that magnetization vanishes and thermodynamical witness (5) reduces to

$$\frac{|U|}{N|J|} > 1. \quad (6)$$

In Ref. [7] it was shown that concurrence C (as a measure of bipartite entanglement [21]) is zero at any temperature in the ferromagnetic case and that it is given by $C = \frac{1}{2} \max [0, |U|/(NJ) - 1]$ in the antiferromagnetic case. Thus C is nonzero if and only if $|U|/(NJ) > 1$. This shows that our thermodynamical entanglement witness can detect entire bipartite entanglement as measured by concurrence. Furthermore, the fact that the value of the entanglement witness for the ground state is well above the limit of 1 ($|E_0/JN| = 1.773$) suggests that entanglement could exist and be detected by the thermodynamical witness at nonzero temperatures as well.

Next we analyze *XXX Heisenberg model with nonzero magnetic field* ($J_x = J_y = J_z = J$ and $B \neq 0$). Recently, it was shown that concurrence vanishes in the thermodynamical limit for the ferromagnetic case ($J < 0$) at low temperatures when only ground state and the first excited states (single spin excitations) are populated [15]. Within the validity of this approximation the partition function is given by [15] $Z =$




FIG. 1: The parameter regions of temperature $kT/|J|$ and external magnetic field $B/|J|$ (expressed in the units of the coupling constant J ; k is the Boltzmann constant) where thermodynamical entanglement witness $|(U+MB)/(NJ)| > 1$ detects entanglement in the XX Heisenberg solid state system both in ferromagnetic and antiferromagnetic case.

$e^{B\beta(J+B)}(1 + e^{-2\beta B}N/\sqrt{8\pi\beta J})$. Using this we obtain $|U + BM|/(NJ) = 1$ and thus no entanglement can be detected at the level of approximation in agreement with the result of Ref. [15].

We proceed with the consideration of *XX Heisenberg model with nonzero magnetic field* ($J_x = J_y = J$, $J_z = 0$ and $B \neq 0$). This case is the most interesting as it is exactly solvable and the partition function was found in Ref. [26]. Let us introduce the following dimensionless quantities: $C = B/kT$ and $K = J/kT$ (note a difference of factor 2 in the definitions of J and K with respect to Ref. [26]) and the function

$$f(K, C, \omega) = \sqrt{2K^2 + 2K^2 \cos 2\omega - 4CK \cos \omega + C^2} \quad (7)$$

for convenience. Then the internal energy is given by [26]

$$\frac{U}{N} = -\frac{kT}{\pi} \int_0^\pi f(K, C, \omega) \tanh f(K, C, \omega) d\omega, \quad (8)$$

and the magnetization by [26]

$$\frac{M}{N} = -\frac{1}{\pi} \int_0^\pi \frac{4K^2 \cos^2 \omega}{f(K, C, \omega)} \tanh f(K, C, \omega) d\omega \quad (9)$$

both in ferromagnetic and antiferromagnetic case.

We use Eqs. (8) and (9) to determine the parameter regions of temperature kT and magnetic field B for which entanglement exists in the solid state system (Fig. 1). The critical values of kT and B below which entanglement can be detected is of the order of J , which can be as high as 10 Kelvin [23].

Finally, we consider *XX Heisenberg model with no magnetic field* ($J_x = J_y = J$, $J_z = 0$ and $B = 0$). This model is important for quantum information processing as it describes the effective interaction in cavity QED and between two quantum dots [27]. It also can be used to

construct the controlled-NOT gate [27]. Since there is no magnetic field, magnetization vanishes and thermodynamical witness reduces again to Eq. (6). Nevertheless, the expressions (8) for internal energy and (9) for magnetization are valid also for the case with no magnetic field. Thus the axes $B = 0$ on Fig. 1 gives the temperature interval within which entanglement is present in the XX system.

The common feature in all cases for which the existence of entanglement could here be proven is that both high temperatures and high values of magnetic field move the thermal states away from the region with non-zero entanglement. This is understandable because high values of magnetic field tend to order all spins parallel to the field which corresponds to an overall state being a product of the individual spin states. Increasing the temperature has also entanglement destructive character due to thermal fluctuations (which, on the other hand, can be understood due to the interaction between the environment and the solid state system).

We note that our method for determining entanglement in solids within the models of Heisenberg interaction is useful in the cases where other methods fail due to incomplete knowledge of the system. This is the case when only the eigenvalues but not eigenstates of the Hamiltonian are known (which is the most usual case in solid state physics) and thus no measure of entanglement can be computed. Furthermore, in the cases where we lack complete description of the systems we can approach the problem experimentally and determine the value of the thermodynamical entanglement witness by performing appropriate measurements.

Our work raises a number of interesting questions and possibilities for generalizations such as consideration of Hamiltonians with higher spins, two- and three-dimensional systems, non-nearest interactions, anisotropies, other thermodynamical properties (e.g. heat capacity, magnetic susceptibility) and so on.

In conclusion, we show that the presence of entanglement in solid state systems at nonzero temperatures in the thermodynamical limit can be detected by measuring solely macroscopic thermodynamical properties. The parameter regions of temperature and magnetic field are determined for which there is entanglement in the systems. Besides being of fundamental interest, we expect our results to be useful for potential physical realization of a future quantum computer. In the opinion of many researchers, if the future computer is supposed to reach the stage of wide commercial application, it should be based on solid states systems. It will thus be important to derive the critical values of physical parameters (e.g. high-temperature limit) above which one can not harness quantum entanglement in solids as a resource for quantum information processing.

Č.B. has been supported by the European Commission, Marie Curie Fellowship, Project No. 500764 and

by the Austrian Science Foundation (FWF) Project No. F1506. V.V. acknowledges EPSRC for financial support.

* Electronic address: caslav.brukner@univie.ac.at

† Electronic address: v.vedral@imperial.ac.uk

- [1] E. Schrödinger, *Naturwissenschaften* **23**, 807-812, 823-828, 844-849, (1935).
- [2] J.S. Bell, *Physics* (Long Island City, N.Y.) **1**, 195 (1964).
- [3] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University, 2000).
- [4] M.A. Nielsen, PhD Thesis, University of New Mexico (1998), e-print quant-ph/0011036.
- [5] M.C. Arnesen, S. Bose and V. Vedral, *Phys. Rev. Lett.* **87**, 017901 (2001).
- [6] D. Gunlycke, S. Bose, V.M. Kendon and V. Vedral, *Phys. Rev. A* **64** 04302 (2001).
- [7] X. Wang and P. Zanardi, *Phys. Lett. A* **301** 1 (2002).
- [8] X. Wang, *Phys. Rev. A* **66** 034302 (2002).
- [9] K.M. O'Connor and W.K. Wootters, *Phys. Rev. A* **63** 052302 (2001).
- [10] T.J. Osborne and M.A. Nielsen, *Phys. Rev. A* **66** 032110 (2002).
- [11] A. Osterloh, L. Amico, G. Falci and R. Fazio, *Nature* **416**, 608 (2002).
- [12] G. Vidal, J. I. Latorre, E. Rico and A. Kitaev, *Phys. Rev. Lett.* **90** (2003) 227902.
- [13] J.I. Latorre, E. Rico and G. Vidal, *Quant. Inf. and Comp.* **4** (2004) 048-092.
- [14] B.-Q. Jin and V.E. Korepin, *Phys. Rev. A* **69**, 120406 (2004).
- [15] M. Asoudeh and V. Karimipour, e-print quant-ph/0405060.
- [16] V. Vedral, *N. J. Phys.* **6**, 22 (2004).
- [17] V. Vedral, e-print quant-ph/0405102.
- [18] H. Fan and S. Lloyd, e-print quant-ph/0405130.
- [19] S. Ghosh, T.F. Rosenbaum, G. Aeppli, S.N. Copper-smith, *Nature* **425**, 48 (2003); V. Vedral, *Nature* **425**, 28 (2003).
- [20] S. Sachdev, *Quantum Phase Transitions* (Cambridge University Press, Cambridge, 1999).
- [21] W.K. Wootters, *Phys. Rev. Lett.* **80**, 2245 (1998).
- [22] M. Horodecki, P. Horodecki and R. Horodecki, *Phys. Lett. A* **223**, 1 (1996); D. Bruss, *J. Math. Phys.* **43**, 4237 (2002); B. M. Terhal, *J. Th. Comp. Sc.* **287**(1), 313 (2002).
- [23] See, for example: P.R. Hammar, M.B. Stone, D.H. Reich, C. Broholm, P.J. Gibson, M.M. Turnbull, C.P. Landee, and M. Oshikawa, *Phys. Rev. B* **59**, 1008 (1999).
- [24] G. Toth, O. Gühne and J.I. Cirac, Poster at Gordon Research Conference on Quantum Information, Ventura, California, February 22-27, 2004.
- [25] L. Hulthén, *Arkiv Mat. Astron. Fysik* **26A**, No 1 (1938).
- [26] S. Katsura, *Phys. Rev.* **127**, 1508 (1962).
- [27] A. Imamoglu, D.D. Awschalom, G. Burkard, D.P. Di Vigenzo, D. Loss, M. Sherwin, and A. Small, *Phys. Rev. Lett.* **83**, 4204 (1999).